

UNIVERSAL TORSORS OVER QUARTIC DEL PEZZO SURFACES AND STABLE RATIONALITY

YURI TSCHINKEL AND ZHIJIA ZHANG

ABSTRACT. Let S be a smooth quartic del Pezzo surface over a field k , of characteristic zero. We prove that a universal torsor $\mathcal{T}$ over S is k -rational, provided $\mathcal{T}$ has k -points. As an application, we obtain examples of stably rational smooth cubic hypersurfaces over $\mathbb{Q}$ in every dimension greater than 2.

1. INTRODUCTION

Let S be a del Pezzo surface over a field k with a nonempty set $S(k)$ of k -rational points. Let

$$\mathcal{T} \rightarrow S$$

be a universal torsor over S , under the Néron-Severi torus of S ; see Section 2 for details on the terminology. Our first result is a proof of a conjecture of Colliot-Thélène–Sansuc [CTS87a, Section 2.8], for quartic del Pezzo surfaces:

Theorem 1.1 (Theorem 3.4). *Let k be a field of characteristic zero and S a quartic del Pezzo surface over k . Let $\mathcal{T} \rightarrow S$ be a universal torsor over S with $\mathcal{T}(k) \neq \emptyset$. Then $\mathcal{T}$ is k -rational.*

Rationality of universal torsors is a key property in the study of the Hasse principle and weak approximation on del Pezzo surfaces over number fields, see, e.g., [CTS87a] and [SS91]. Here, we obtain applications to stable rationality:

Corollary 1.2 (Corollary 4.3). *Let k be a field of characteristic zero and S a quartic del Pezzo surface over k . Then S is stably rational over k if and only if*

- $S(k) \neq \emptyset$ and
- S satisfies Condition **(H1)**, i.e., vanishing of Galois cohomology

$$H^1(\mathfrak{g}_{k'}, \text{Pic}(\bar{S})) = 0, \quad \text{for all finite } k'/k. \quad (1.1)$$

Date: August 26, 2026.

2020 Mathematics Subject Classification. 14E08; 14E05, 14M20.

Rationality of del Pezzo surfaces is well-understood in terms of the Galois action on the geometric Picard group $\text{Pic}(\bar{S})$, see, e.g., [Man86]. For quartic del Pezzo surfaces S , the Galois action factors through a subgroup of $W(\mathbf{D}_5)$, the Weyl group of the root system $\mathbf{D}_5$. There are 4 types of Galois actions that correspond to nonrational S satisfying the conditions of Corollary 1.2, see [KiST89, Theorems 4.19 and 5.20]. Explicit presentations of these actions can be found in Section 4 (see also [TY20]). We list them by their images in $W(\mathbf{D}_5)$:

- $I_0 - \mathfrak{S}_3$,
- $I_1 - \mathfrak{C}_2 \times \mathfrak{S}_3$,
- $I_2 - \mathfrak{C}_3 \rtimes \mathfrak{C}_4$,
- $I_3 - \mathfrak{C}_3 \rtimes \mathfrak{D}_4$.

Rationality of $\mathcal{T}$ in the case I_0 was established in [BCTSSD85, Section 1]. The corresponding S admits a conic bundle over k , and $\text{rkPic}(S) = 2$. The main application in [BCTSSD85] produced the first example of a stably rational but nonrational threefold X , as a smooth projective model of a quartic del Pezzo surface over the function field $k = \mathbb{C}(t)$, with Galois action of type I_0 . Stable rationality of X over $\mathbb{C}$ follows formally from the stable rationality of S over k . On the other hand, X is also a conic bundle over a rational surface, and the intermediate Jacobian of X is the Prym variety of the discriminant curve, with the associated double cover. Choosing an appropriate configuration of the discriminant curve, one can ensure that X is nonrational.

This beautiful construction served as input for [HKT23] to produce smooth families of threefolds with stably rational and stably nonrational fibers. It seemed natural to try to apply it to other nonrational threefolds, e.g., cubic threefolds. This was sketched in unpublished notes [HT17a], [HT17b], where it was realized that:

- the I_0 case cannot arise from a smooth cubic threefold (birationally) as a quartic del Pezzo surface fibration,
- a proof of stable rationality in the I_1 case over $k = \mathbb{C}(t)$ would produce stably rational smooth cubic threefolds over $\mathbb{C}$.

The bottleneck that remained was a proof of rationality of the torsor. We settle this in Section 3. In Section 4, we apply this to a proof of Corollary 1.2. The proof follows exactly the strategy outlined in [BCTSSD85, Section 1]:

- stable rationality of the Néron-Severi torus T of S – this follows from the fact that, in our situation, $\text{Pic}(\bar{S})$ is a stably permutation module,
- existence of a universal torsor $\mathcal{T}$ over S with $\mathcal{T}(k) \neq \emptyset$, and k -rationality of $\mathcal{T}$,

- k -birationality of $\mathcal{T}$ to $S \times T$, and thus stable rationality of S over k .

In Section 5, we prove:

Theorem 1.3 (Propositions 5.1 and 5.3). *For every $n \geq 3$ there exist stably rational smooth cubic hypersurfaces $X \subset \mathbb{P}^{n+1}$, over $\mathbb{Q}$.*

Combining Theorem 1.3 with the recent proof in [EFS25] that a very general cubic threefold over $\mathbb{C}$ does not admit a decomposition of the diagonal, and is thus not stably rational over $\mathbb{C}$, we obtain new smooth families of threefolds with stably rational and stably nonrational fibers, extending the results of [HKT23].

The proof of Theorem 1.1 was arrived at via a detour through equivariant geometry, that provided some rigidity in the quest for a rationality construction.

The analogies between geometry over nonclosed fields and equivariant geometry have been discussed elsewhere, e.g., [KT26a], [KT26b]. In particular, there is also a Condition **(H1)**, which is necessary for stable linearizability of a regular action of a finite group G on a del Pezzo surface over an algebraically closed field, identical to (1.1), but with Galois cohomology replaced by group cohomology:

$$H^1(G', \text{Pic}(S)) = 0, \quad \forall G' \subseteq G.$$

In the situation at hand, only the I_2 -action can be realized as a geometric action on a del Pezzo surface over an algebraically closed field of characteristic zero, by [Pro15, Theorem 1.2]. The following answers the last remaining open question in this direction:

Theorem 1.4. *Let S be the quartic del Pezzo surface given by*

$$S = \{x_1^2 + \zeta_3 x_2^2 + \zeta_3^2 x_3^2 + x_4^2 = x_1^2 + \zeta_3^2 x_2^2 + \zeta_3 x_3^2 + x_5^2 = 0\} \subset \mathbb{P}^4,$$

with an action of $G = \mathfrak{C}_3 \rtimes \mathfrak{C}_4$ generated by

$$\begin{aligned} (x_1, \dots, x_5) &\mapsto (x_2, x_3, x_1, \zeta_3 x_4, \zeta_3^2 x_5), \\ (x_1, \dots, x_5) &\mapsto (x_1, x_3, x_2, -x_5, x_4). \end{aligned}$$

Then the G -action on S is stably linearizable.

We prove this in Section 6, by writing down one more time a suitable universal torsor $\mathcal{T}$, lifting the G -action to $\mathcal{T}$, and proving the linearizability of $\mathcal{T}$. The key observation is that $\mathcal{T}$ has a G -fixed point. This is the first nontrivial example of a stably linearizable but not coarsely linearizable action on a rational variety, in the terminology of [KT26b].

Acknowledgments: We are grateful to A. Kresch and B. Hassett for many years of collaboration on this and related problems, and to J.-L. Colliot-Thélène for his interest and suggestions. We used Claude Fable 5 and ChatGPT-5.6 Sol for searching the literature, testing hypotheses, checking computations, and proofreading. The paper was written by the authors, who are responsible for its correctness.

2. GENERALITIES

The material in this section is standard, and is assembled here for convenience. Throughout, k is a field of characteristic zero. We fix an algebraic closure $\bar{k}$ of k and let $\mathfrak{g} = \mathfrak{g}_k$ be the Galois group of $\bar{k}/k$. For a variety X over k , we let $\bar{X}$ be the base change to $\bar{k}$.

Modules. Let G be a profinite group, with continuous action on a discrete finite-rank $\mathbb{Z}$ -module M , and

$$H^1(G, M)$$

the first cohomology group of M . A G -module M is called a *permutation module* if it has a $\mathbb{Z}$ -basis permuted by G , and a *stably permutation module* if there exist permutation G -modules P and P' , and an isomorphism of G -modules $M \oplus P \simeq P'$. By Shapiro's lemma, if M is stably permutation, then $H^1(H, M) = 0$, for every open subgroup $H \subseteq G$.

Tori. There is a duality between the category of algebraic tori over k and discrete continuous finite-rank $\mathfrak{g}$ -modules, via the assignment

$$T \mapsto \mathfrak{X}^*(\bar{T}),$$

sending T to the group of characters of $\bar{T}$. Arithmetic properties of T are reflected in properties of the $\mathfrak{g}$ -module $\mathfrak{X}^*(\bar{T})$, e.g., if $\mathfrak{X}^*(\bar{T})$ is a stably permutation $\mathfrak{g}$ -module, then T is stably rational over k [CTS77, Section 2].

Universal torsors. Let T be an algebraic torus over k and S a smooth projective variety over k , with $\text{Pic}(\bar{S})$ torsion-free and of finite rank, e.g., a del Pezzo surface. There is a natural homomorphism

$$H^1(S, T) \xrightarrow{\chi} \text{Hom}_{\mathfrak{g}}(\mathfrak{X}^*(\bar{T}), \text{Pic}(\bar{S}))$$

from the group classifying isomorphism classes of T -torsors over S , see [CTS87a, Theorem 1.5.1]. When T is the Néron-Severi torus, i.e.,

$$\mathfrak{X}^*(\bar{T}) \simeq \text{Pic}(\bar{S}),$$

a T -torsor $\mathcal{T}$ is called *universal* if

$$\chi([\mathcal{T}]) = \text{Id}.$$

Such torsors exist over $\bar{k}$ but not always over k . However, they do exist under the assumption that $S(k) \neq \emptyset$ [CTS87a, Remark 2.2.9].

Lemma 2.1. *Let S be a del Pezzo surface of degree ≥ 3 over k and $\mathcal{T}$ a universal torsor over S . If $\mathcal{T}(k) \neq \emptyset$ then $\mathcal{T}(k)$ is Zariski dense in $\mathcal{T}$.*

Proof. Let $\xi : \mathcal{T} \rightarrow S$ be the torsor and $t \in \mathcal{T}(k)$. Then $S(k)$ is nonempty, as it contains $s := \xi(t)$, and S is k -unirational, by [Man86, Theorems 29.4 and 30.1]. In degree ≥ 6 , universal torsors are Zariski open subsets of affine space. In degree 5, a universal torsor embeds as a dense Zariski open subset into the affine cone over the Grassmannian $\mathrm{Gr}(2, 5)$, and rational points on it are dense as well.

It remains to treat degree 3 and 4. The specialization $\mathcal{T}_s$ of $\mathcal{T}$ to s is split, i.e., the class

$$[\mathcal{T}_s] \in H^1(\mathrm{Spec}(k), T)$$

is trivial, since there is a point $t \in \mathcal{T}_s(k)$. This implies that the fiber is isomorphic to T , and has Zariski dense k -rational points.

By [CTS87a, Proposition 2.7.2], specializations of the universal torsor $\mathcal{T}$ to R -equivalent points in $S(k)$ have the same class in $H^1(\mathrm{Spec}(k), T)$. It suffices to show that points in $S(k)$ which are R -equivalent to s are Zariski dense in S . Indeed, for each $s' \in S(k)$ in the same R -equivalence class as s , the torsor $\mathcal{T}_{s'}$ is split. Each such fiber has a Zariski dense set of rational points. Thus k -points are Zariski dense in $\mathcal{T}$.

For smooth cubic surfaces, this follows from [Man86, Theorems 13.1 and 14.3]. The case of quartic del Pezzo surfaces S can be reduced to the cubic surface case via a blowup of a general k -point (distinct from s and away from exceptional curves), or handled via a unirational parametrization $\mu : \mathbb{P}^2 \dashrightarrow S$ containing s in its image: all points in the image of μ will be in the same R -equivalence class and Zariski dense. Arguments of this type can be found in [CTS87a, Section 2.9]. $\square$

Remark 2.2. In our applications to stable rationality, the character lattice $\mathfrak{X}^*(\bar{T})$ is a stably permutation $\mathfrak{g}$ -module, with

$$H^1(\mathrm{Spec}(K), T) = 0,$$

for any field K/k , by Hilbert's Theorem 90. In particular, the T -torsor $\mathcal{T} \rightarrow S$ splits over the function field $k(S)$, and every fiber over a point $s \in S(k)$ is isomorphic to T , over k .

Stable rationality. Recall that an algebraic variety X is rational if X is birational to $\mathbb{P}^n$, and is stably rational if $X \times \mathbb{P}^r$ is rational, for some integer r . The application of the theory of universal torsors to stable rationality was pioneered in [BCTSSD85], which led to the first (and

so far only) examples of stably rational but nonrational varieties. The stable rationality construction relies on the following:

Proposition 2.3 ([BCTSSD85, Proposition 3]). *Let X be a smooth, proper, geometrically integral, and geometrically rational variety over k . Assume that there exists a universal torsor $\mathcal{T} \rightarrow X$ which is a k -rational variety, and the $\mathfrak{g}$ -module $\mathrm{Pic}(\bar{X})$ is a stably permutation module. Then X is stably rational over k .*

The proof of Proposition 2.3 uses Hilbert's Theorem 90 and Shapiro's lemma to establish a birationality

$$\mathcal{T} \sim_k X \times T,$$

where T is the Néron-Severi torus of X . Stable rationality of X then follows from the assumptions on $\mathcal{T}$ and $\mathrm{Pic}(\bar{X})$.

OADP varieties. Recall that an n -dimensional reduced, irreducible, projective variety $X \subset \mathbb{P}^{2n+1}$ has *one apparent double point* if through a general point in $\mathbb{P}^{2n+1}$, there passes a unique secant line of X . See [CR06] for a discussion of properties of such varieties; rationality constructions based on apparent double points can be found in, e.g., [CMR04], [HKT22, Sections 3 and 7]. Our proof of rationality of universal torsors $\mathcal{T}$ over quartic del Pezzo surfaces will be based on the following strengthening of [CR06, Theorem 2.7] (see also, [CMR04, Theorem 4.1]):

Theorem 2.4. *Let $X \subset \mathbb{P}^{2n+1}$ be an n -dimensional reduced, irreducible, nondegenerate projective variety over a field k of characteristic zero. Assume that $X(k)$ is Zariski dense and that $\bar{X}$ has one apparent double point. Then the projection*

$$X \dashrightarrow \mathbb{P}^n$$

from the tangent space of a general point $x \in X(k)$ is a birational map.

Proof. The statement in [CR06, Theorem 2.7] is formulated over $\mathbb{C}$. But the proof works for any algebraically closed field $\bar{k}$ of characteristic 0. The extension to a nonclosed field k is immediate: the projection map from the tangent space of a general k -point is defined over k , and is birational over $\bar{k}$. Thus it is birational over k . Zariski density of $X(k)$ allows to pick an appropriate point. $\square$

3. UNIVERSAL TORSORS OVER QUARTIC DEL PEZZO SURFACES

We recall the construction of universal torsors over quartic del Pezzo surfaces. In the geometric case, this has been worked out in [Der07], [BP04], [SS07]. For results over nonclosed fields, see [CTS87a], [DP19].

We start with the geometric case, i.e., when $k = \bar{k}$. A universal torsor admits an embedding

$$\mathcal{T} \hookrightarrow \text{Spec}(\text{Cox}(S)),$$

as a dense Zariski open subset; here $\text{Cox}(S)$ is the *Cox ring* of a quartic del Pezzo surface S . Concretely, let $S \rightarrow \mathbb{P}^2$ be given as a blowup of 5 points in general position, which can be chosen to be

$$p_1 = [1 : 0 : 0], \quad p_2 = [0 : 1 : 0], \quad p_3 = [0 : 0 : 1], \quad p_4 = [1 : 1 : 1],$$

$$p_5 = [1 : a : b],$$

where $a, b \in k$ satisfy

$$ab(a-1)(b-1)(a-b) \neq 0.$$

The Cox ring $\text{Cox}(S)$ is generated by 16 sections corresponding to exceptional curves in S modulo 20 relations arising from 10 rulings in S , see [BP04, Theorem 4.9]. The 16 exceptional curves are

- 5 exceptional curves E_i above p_i ,
- 10 strict transforms L_{ij} of lines passing through p_i and p_j ,
- the strict transform Q of the conic passing through the 5 points.

The corresponding generators of $\text{Cox}(S)$ are denoted by

$$e_i, \quad l_{ij}, \quad q.$$

To each exceptional curve we assign a form defining its image in $\mathbb{P}_{z_1, z_2, z_3}^2$

$$f_{E_i} = 1, \quad f_{L_{12}} = z_3, \quad f_{L_{13}} = z_2, \quad f_{L_{14}} = z_2 - z_3, \quad f_{L_{15}} = bz_2 - az_3,$$

$$f_{L_{23}} = z_1, \quad f_{L_{24}} = z_1 - z_3, \quad f_{L_{25}} = bz_1 - z_3, \quad f_{L_{34}} = z_1 - z_2,$$

$$f_{L_{35}} = az_1 - z_2, \quad f_{L_{45}} = (b-a)z_1 + (1-b)z_2 + (a-1)z_3,$$

$$f_Q = b(1-a)z_1z_2 + a(b-1)z_1z_3 + (a-b)z_2z_3.$$

These determine relations in $\text{Cox}(S)$. The 10 rulings in S are classes

$$H - E_i, \quad 2H + E_i - \sum_{j=1}^5 E_j, \quad i = 1, \dots, 5,$$

where H is the pullback of a general line on $\mathbb{P}^2$. We compute the relations associated with each ruling: For each $i = 1, \dots, 5$, there are 4 reducible members in the pencil $|H - E_i|$,

$$L_{ij} + E_j, \quad j \in \{1, \dots, 5\} \setminus \{i\},$$

yielding relations given by the vanishing of the following forms:

$$\begin{aligned}
R_1 &= e_2 l_{12} - e_3 l_{13} + e_4 l_{14}, \\
R_2 &= a e_2 l_{12} - b e_3 l_{13} + e_5 l_{15}, \\
R_3 &= e_1 l_{12} - e_3 l_{23} + e_4 l_{24}, \\
R_4 &= e_1 l_{12} - b e_3 l_{23} + e_5 l_{25}, \\
R_5 &= e_1 l_{13} - e_2 l_{23} + e_4 l_{34}, \\
R_6 &= e_1 l_{13} - a e_2 l_{23} + e_5 l_{35}, \\
R_7 &= e_1 l_{14} - e_2 l_{24} + e_3 l_{34}, \\
R_8 &= (b-1) e_1 l_{14} + (a-b) e_2 l_{24} + e_5 l_{45}, \\
R_9 &= e_1 l_{15} - a e_2 l_{25} + b e_3 l_{35}, \\
R_{10} &= (a-1) e_2 l_{25} + (1-b) e_3 l_{35} + e_4 l_{45}.
\end{aligned}$$

For $i = 1, \dots, 5$, the 4 reducible members in $|2H + E_i - \sum_{j=1}^5 E_j|$ are

$$Q + E_i, \quad L_{i_1 i_2} + L_{i_3 i_4}, \quad \{i_1, i_2, i_3, i_4\} = \{1, \dots, 5\} \setminus \{i\}.$$

The remaining relations are given by the vanishing of the following forms:

$$\begin{aligned}
R_{11} &= l_{23} l_{45} + l_{24} l_{35} - l_{25} l_{34}, \\
R_{12} &= a l_{23} l_{45} + (a-b) l_{24} l_{35} - e_1 q, \\
R_{13} &= l_{13} l_{45} + l_{14} l_{35} - l_{15} l_{34}, \\
R_{14} &= l_{13} l_{45} + (1-b) l_{14} l_{35} - e_2 q, \\
R_{15} &= l_{12} l_{45} + l_{14} l_{25} - l_{15} l_{24}, \\
R_{16} &= l_{12} l_{45} + (1-a) l_{14} l_{25} - e_3 q, \\
R_{17} &= l_{12} l_{35} - l_{13} l_{25} + l_{15} l_{23}, \\
R_{18} &= (b-1) l_{12} l_{35} + (1-a) l_{13} l_{25} - e_4 q, \\
R_{19} &= l_{12} l_{34} - l_{13} l_{24} + l_{14} l_{23}, \\
R_{20} &= a(b-1) l_{12} l_{34} + b(1-a) l_{13} l_{24} - e_5 q.
\end{aligned}$$

Then

$$\text{Cox}(S) = k[q, e_1, \dots, e_5, l_{ij}] / \langle R_1, \dots, R_{20} \rangle, \quad 1 \leq i < j \leq 5,$$

and

$$U := \text{Spec}(\text{Cox}(S)) \subset \mathbb{A}^{16} \tag{3.1}$$

contains a universal torsor $\mathcal{T}$ over S as a dense Zariski open subset. Taking the projectivization gives $\mathbb{P}(U) \subset \mathbb{P}^{15}$. Let

$$J := \left(\frac{\partial R_i}{\partial x_j} \right) \tag{3.2}$$

be the Jacobian matrix, where x_j is the j th generator of $\text{Cox}(S)$, with the natural lexicographic order on l_{ij} .

Lemma 3.1. *Let u_1, u_2 be two points on U . Then*

$$u_1 - u_2 \in \ker(J(u_1 + u_2)).$$

Proof. Let $B_i(x, y) = R_i(x + y) - R_i(x) - R_i(y)$ be the symmetric bilinear form associated with the quadratic form R_i . For any $t \in k$, we have that

$$R_i(x + ty) = tB_i(x, y) + R_i(x) + t^2R_i(y).$$

The i th component of $J(x) \cdot y$ is

$$(J(x) \cdot y)_i = \left. \frac{d}{dt} \right|_{t=0} R_i(x + ty) = B_i(x, y).$$

It follows that for every $i = 1, \dots, 20$,

$$(J(u_1 + u_2) \cdot (u_1 - u_2))_i = B_i(u_1 + u_2, u_1 - u_2) = 2R_i(u_1) - 2R_i(u_2) = 0.$$

□

The key observation is that $\mathbb{P}(U)$ is an OADP variety:

Lemma 3.2. *Through a general point in $\mathbb{P}^{15}$, there passes a unique secant line of $\mathbb{P}(U)$.*

Proof. First, we show that the secant variety of $\mathbb{P}(U)$ is the whole $\mathbb{P}^{15}$. By Terracini's Lemma (see, e.g., [CR06, Theorem 1.1]), it suffices to show that for two general points $u_1, u_2 \in U$, we have that

$$T_{u_1}(U) + T_{u_2}(U) = \mathbb{A}^{16}, \quad (3.3)$$

where $T_{u_i}(U)$ is the embedded tangent space of U at u_i , $i = 1, 2$. Consider

$$t_1 = (0, 1, 1, 0, \dots, 0), \quad t_2 = (1, 0, 0, 0, 0, 0, 1, 0, \dots, 0).$$

A computation¹ of the Jacobian matrix J defined in (3.2) shows that t_1, t_2 are smooth points of U and $T_{t_1}(U) + T_{t_2}(U) = \mathbb{A}^{16}$. Since (3.3) is an open condition, it holds for two general points on U .

Let $x \in \mathbb{A}^{16}$ be a general point. We now show that there is exactly one secant line of $\mathbb{P}(U)$ passing through the image of x in $\mathbb{P}^{15}$. Assume that two such lines l_1 and l_2 exist. We can find points t_3, t_4, t'_3, t'_4 on U such that $l_1 = \mathbb{P}(\langle t_3, t_4 \rangle)$, $l_2 = \mathbb{P}(\langle t'_3, t'_4 \rangle)$, and $x = t_3 + t_4 = t'_3 + t'_4$. By Lemma 3.1, we know that

$$J(x) \cdot (t_3 - t_4) = J(x) \cdot (t'_3 - t'_4) = 0.$$

A direct computation shows that $\dim(\ker(J(t_1 + t_2))) = 1$, which implies that $\dim(\ker(J(x))) \leq 1$, since this is an open condition and x is general. It follows that $t_3 - t_4$ and $t'_3 - t'_4$ are linearly dependent, i.e., $l_1 = l_2$. □

¹See [TZ26] for supporting materials.

Remark 3.3. In any concrete situation, one can directly check the OADP property of $\mathbb{P}(U)$ and the birationality of the projection in Theorem 2.4. An example of this computation is provided in Section 6.

Theorem 3.4. *Let S be a quartic del Pezzo surface over k and $\mathcal{T} \rightarrow S$ a universal torsor over S . If $\mathcal{T}(k) \neq \emptyset$, then $\mathcal{T}$ is rational over k .*

Proof. By Lemma 2.1, $\mathcal{T}(k)$ is Zariski dense in $\mathcal{T}$. The $\mathfrak{g}$ -equivariant structures

$$\bar{\mathcal{T}} \hookrightarrow \bar{U} := \operatorname{Spec}(\operatorname{Cox}(\bar{S})) \subset \mathbb{A}^{16} \quad (3.4)$$

descend to k , i.e., we have an embedding

$$\mathcal{T} \hookrightarrow U \subset \mathbb{A}^{16}, \quad (3.5)$$

over k , such that (3.4) is the base change of (3.5) to $\bar{k}$, see, e.g., [DP19, Remark 4.5]. On passage to $\bar{k}$, the universal torsor $\bar{\mathcal{T}}$ with the embedding (3.4) is linearly isomorphic to the embedding of the universal torsor constructed in (3.1). Geometrically, $\bar{\mathcal{T}}$ is a Zariski dense open subset of a cone over a variety $\mathbb{P}(\bar{U})$ which has one apparent double point by Lemma 3.2. Note that $\mathbb{P}(\bar{U})$ is reduced, irreducible and nondegenerate, since $\operatorname{Cox}(\bar{S})$ is an integral domain and there are no linear forms in the defining ideal. By Theorem 2.4, projection from the tangent space of a general k -point in $\mathbb{P}(U)$ yields k -rationality of $\mathbb{P}(U)$ and thus of $\mathcal{T}$. $\square$

Corollary 3.5. *Let S be a quartic del Pezzo surface over k such that $S(k) \neq \emptyset$. Then there exists a k -rational universal torsor $\mathcal{T} \rightarrow S$.*

Proof. By Theorem 3.4, it suffices to show the existence of a universal torsor $\mathcal{T} \rightarrow S$ with a rational point. Recall that $S(k) \neq \emptyset$ implies that we can find $s \in S(k)$ in the complement to exceptional curves. By [CTS87a, Corollary 2.3.4], there exists a universal torsor $\mathcal{T}$ over S whose fiber over s is a trivial torsor, and thus $\mathcal{T}(k) \neq \emptyset$. $\square$

4. STABLE RATIONALITY OF QUARTIC DEL PEZZO SURFACES

We recall basic facts about the Picard group of a quartic del Pezzo surface S , see [Man86] and [KiST89]. The anticanonical model of S is a smooth intersection of two quadrics

$$S = Q_1 \cap Q_2 \subset \mathbb{P}^4.$$

The pencil of quadrics generated by Q_1 and Q_2 has 5 singular members, over $\bar{k}$. Each of them is a cone over $\mathbb{P}^1 \times \mathbb{P}^1$, whose two rulings give rise to two conic bundle structures on $\bar{S}$. Choosing an appropriate contraction

$\bar{S} \rightarrow \mathbb{P}^2$ and labeling the 16 exceptional curves in $\bar{S}$ by Q, E_i, L_{ij} as in Section 3, the classes of 10 rulings in $\text{Pic}(\bar{S})$ are

$$F_i := H - E_i, \quad F'_i = 2H + E_i - \sum_{j=1}^5 E_j, \quad i = 1, \dots, 5, \quad (4.1)$$

where H is the pullback of a general line on $\mathbb{P}^2$. Each pair $\{F_i, F'_i\}$ consists of the two rulings of the same singular quadric.

The $\mathfrak{g}$ -action on $\text{Pic}(\bar{S})$ factors through a subgroup of the Weyl group

$$W(D_5) \simeq \mathfrak{C}_2^4 \rtimes \mathfrak{S}_5,$$

where $\mathfrak{S}_5$ permutes the 5 singular members of the pencil, and $\mathfrak{C}_2^4$ is generated by elements simultaneously switching the rulings of the i th and j th singular members, denoted by $c_i c_j$.

Choose a bijection between the 16 exceptional curves, and the 16 odd subsets of $\{1, \dots, 5\}$ given by

$$Q \leftrightarrow \{1, \dots, 5\}, \quad L_{ij} \leftrightarrow \{1, \dots, 5\} \setminus \{i, j\}, \quad E_i \leftrightarrow \{i\}.$$

The $W(D_5)$ -action on exceptional curves can be described in terms of its action on odd subsets of $\{1, \dots, 5\}$ as follows: $\mathfrak{S}_5$ permutes the 5 indices, and $c_i c_j$ changes an odd subset I to J where for any $r \in \{1, \dots, 5\}$,

- if $r \in I$, then $r \in J$ if and only if $r \notin \{i, j\}$,
- if $r \notin I$, then $r \in J$ if and only if $r \in \{i, j\}$.

Here is an alternative way to identify the image of $\mathfrak{g}$ in $W(D_5)$, using the blowup $\tilde{S} \rightarrow S$ of a general point $s \in S(k)$ (this will be used in Section 5). Note that $\tilde{S}$ is a cubic surface with a line l above s . We denote the class of l in the geometric Picard group of $\tilde{S}$ by E_6 . The residual conic bundle

$$\varphi : \tilde{S} \rightarrow \mathbb{P}^1 \quad (4.2)$$

has, geometrically, 5 singular fibers, each consisting of two distinct lines. Their components are the 10 lines intersecting l . The i th singular fiber consists of two lines whose classes are

$$H - E_i - E_6, \quad 2H + E_i - \sum_{j=1}^6 E_j, \quad i = 1, \dots, 5.$$

Their images in S are exactly the 10 rulings in (4.1). Therefore, the image of $\mathfrak{g} \rightarrow W(D_5)$ is also determined by the permutation of 5 singular fibers of φ and an even number of switches of their components.

In [Man66], Manin shows that for a stably rational surface S over a field k of characteristic zero, the $\mathfrak{g}$ -module $\text{Pic}(\bar{S})$ is stably permutation. We are interested in the converse.

Proposition 4.1 ([KiST89], [TY20]). *If a quartic del Pezzo surface S is k -minimal, and satisfies Condition **(H1)** of (1.1), i.e.,*

$$H^1(\mathfrak{g}_{k'}, \text{Pic}(\bar{S})) = 0, \quad \text{for all finite } k'/k,$$

then the image of $\mathfrak{g}_k$ in $W(\mathbf{D}_5)$ is conjugate to one of the following:

$$\begin{aligned} (I_0) : \mathfrak{S}_3 &= \langle (3, 4, 5), c_1 c_3 c_4 c_5(4, 5) \rangle, \\ (I_1) : \mathfrak{C}_2 \times \mathfrak{S}_3 &= \langle c_2 c_3(4, 5), c_1 c_2 c_3 c_4(3, 4, 5) \rangle, \\ (I_2) : \mathfrak{C}_3 \rtimes \mathfrak{C}_4 &= \langle (1, 2, 3), c_1 c_2 c_3 c_4(2, 3)(4, 5) \rangle, \\ (I_3) : \mathfrak{C}_3 \rtimes \mathfrak{D}_4 &= \langle c_1 c_2 c_3 c_5(2, 5), c_3 c_4(3, 4)(1, 5, 2) \rangle. \end{aligned}$$

Our presentation of I_1, I_2 , and I_3 is different from that in [KiST89, Theorem 5.20], but the resulting subgroups are conjugate in $W(\mathbf{D}_5)$.

Lemma 4.2. *Let S be a quartic del Pezzo surface over k . Then the $\mathfrak{g}$ -module $\text{Pic}(\bar{S})$ is stably permutation if and only if Condition **(H1)** of (1.1) is satisfied.*

Proof. If S is not k -minimal, the assertion follows from [Man86]. Thus we may assume that S is k -minimal. It suffices to show that $\text{Pic}(\bar{S})$ is stably permutation, for the four types in Proposition 4.1. The type I_0 is addressed in [BCTSSD85]. Up to conjugation in $W(\mathbf{D}_5)$, the groups of type I_0, I_1 and I_2 are subgroups of the group of type I_3 . Thus we only need to consider this last type. Let

$$g_1 = c_1 c_2 c_3 c_5(2, 5), \quad g_2 = c_3 c_4(3, 4)(1, 5, 2).$$

In the standard basis $H, E_1, \dots, E_5$ of $\text{Pic}(\bar{S})$, they act by matrices

$$g_1 = \begin{pmatrix} 3 & 1 & 1 & 1 & 2 & 1 \\ -1 & -1 & 0 & 0 & -1 & 0 \\ -1 & 0 & 0 & 0 & -1 & -1 \\ -1 & 0 & 0 & -1 & -1 & 0 \\ -2 & -1 & -1 & -1 & -1 & -1 \\ -1 & 0 & -1 & 0 & -1 & 0 \end{pmatrix}, \quad g_2 = \begin{pmatrix} 2 & 1 & 1 & 0 & 0 & 1 \\ -1 & -1 & 0 & 0 & 0 & -1 \\ -1 & -1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & -1 & 0 & 0 & -1 \end{pmatrix}.$$

In our convention, the matrices act from the right, e.g.,

$$g_2(H) = 2H - E_1 - E_2 - E_5.$$

We introduce new variables w_0, w_1, w_2, q_0, q_1 with $\mathfrak{g}$ -permutation actions generated by

$$\begin{aligned} g_1(w_0) &= w_0, & g_1(w_1) &= w_2, & g_1(w_2) &= w_1, & g_1(q_0) &= q_1, & g_1(q_1) &= q_0, \\ g_2(w_0) &= w_2, & g_2(w_1) &= w_0, & g_2(w_2) &= w_1, & g_2(q_0) &= q_0, & g_2(q_1) &= q_1. \end{aligned}$$

One can check that

$$\text{Pic}(\bar{S}) \oplus \mathbb{Z}[w_0, w_1, w_2, q_0, q_1]$$

is a permutation module. Indeed, in the basis

$$\begin{aligned}
 b_1 &= 2H - E_2 - E_3 - E_4 - E_5 - w_0 - w_2 - 2q_0 - q_1, \\
 b_2 &= H - E_1 - w_0 - w_1 - q_0 - 2q_1, \\
 b_3 &= 2H - E_1 - E_3 - E_4 - E_5 - w_0 - w_1 - 2q_0 - q_1, \\
 b_4 &= H - E_2 - w_1 - w_2 - q_0 - 2q_1, \\
 b_5 &= H - E_5 - w_0 - w_2 - q_0 - 2q_1, \\
 b_6 &= 2H - E_1 - E_2 - E_3 - E_4 - w_1 - w_2 - 2q_0 - q_1, \\
 b_7 &= -3H + E_1 + E_2 + 2E_3 + E_4 + E_5 + w_0 + w_1 + w_2 + 3q_0 + q_1, \\
 b_8 &= -2H + E_1 + E_2 + E_5 + w_0 + w_1 + w_2 + q_0 + 3q_1, \\
 b_9 &= -H + w_0 + w_1 + w_2 + q_0 + 3q_1, \\
 b_{10} &= -3H + E_1 + E_2 + E_3 + 2E_4 + E_5 + w_0 + w_1 + w_2 + 3q_0 + q_1, \\
 b_{11} &= 3H - E_1 - E_2 - E_3 - E_4 - E_5 - w_0 - w_1 - w_2 - 2q_0 - 2q_1,
 \end{aligned}$$

the permutation action of $\mathfrak{g}$ is visible by expressing

$$\begin{aligned}
 g_1 &= (b_1, b_2)(b_3, b_5)(b_4, b_6)(b_7, b_8)(b_9, b_{10}), \\
 g_2 &= (b_1, b_6, b_3)(b_2, b_5, b_4)(b_8, b_9).
 \end{aligned}$$

□

Corollary 4.3. *Let S be a quartic del Pezzo surface over a field k of characteristic zero such that $S(k) \neq \emptyset$. Then S is stably rational over k if and only if one of the following equivalent conditions holds:*

- (1) $\text{Pic}(\bar{S})$ is a stably permutation $\mathfrak{g}$ -module,
- (2) S satisfies Condition **(H1)** of (1.1),
- (3) S is not k -minimal or the $\mathfrak{g}$ -action on $\text{Pic}(\bar{S})$ factors through one of the four subgroups in Proposition 4.1.

Proof. The equivalence of the conditions follows from Proposition 4.1 and Lemma 4.2. Corollary 3.5 and Proposition 2.3 show that (1) implies the stable rationality of S . The converse is in [Man66]. □

Remark 4.4. Examples of quartic del Pezzo surfaces over $\mathbb{Q}$ of type I_1, I_2 , or I_3 are given in [KiST89, Theorem 6.15]; Corollary 4.3 yields new stably rational but nonrational surfaces over nonclosed fields.

Levels of stable rationality. Let S be a stably rational but nonrational quartic del Pezzo surface. What is the smallest integer r such that $S \times \mathbb{A}^r$ is rational? Our construction of $\text{Pic}(\bar{S})$ as a stably permutation module in Lemma 4.2 gives a uniform upper bound: $S \times \mathbb{A}^{11}$ is rational. One can check that 5 and 11 are the smallest dimensions of I_3 -permutation modules $\mathbf{P}$ and $\mathbf{P}'$ such that $\text{Pic}(\bar{S}) \oplus \mathbf{P} \simeq \mathbf{P}'$. For S of

type I_0 , [BCTSSD85] shows that $r \leq 3$ by working with a torsor under a 3-dimensional torus, which was improved to $r \leq 2$ in [SB04]. It remains open to construct a nonrational variety X such that $X \times \mathbb{A}^1$ is rational.

5. CUBIC HYPERSURFACES

In this section, we provide examples of stably rational smooth cubic hypersurfaces over $\mathbb{Q}$ in every dimension ≥ 3 . No such examples in odd dimensions were previously known. Over $\mathbb{C}$, smooth cubic threefolds are nonrational, by [CG72]. By [EFS25], a very general cubic threefold over $\mathbb{C}$ is not stably rational. After [Voi17], one might expect that stably rational cubic threefolds are dense in moduli – we are unable to reach this conclusion with our constructions.

Let $X \subset \mathbb{P}^{n+1}$ be a smooth cubic hypersurface with $n \geq 3$, and l a line contained in X . Choose a general plane $\Pi \simeq \mathbb{P}^2 \subset \mathbb{P}^{n+1}$ containing l and consider the projection from Π :

$$X \dashrightarrow \mathbb{P}^{n-2}.$$

The generic fiber X_η of the projection is a cubic surface over $K = k(\mathbb{P}^{n-2})$ containing a line l_η defined over K . Contracting l_η yields a quartic del Pezzo surface S birational to X_η over K , with a K -point.

Our goal is to find appropriate X, l , and Π such that the $\mathfrak{g}_K$ -action on $\text{Pic}(S)$ has one of the types I_1 , I_2 , or I_3 . For these, Corollary 4.3 shows that S is stably rational over K , and thus X is stably rational over k .

Type I_1 . Here, we construct a smooth cubic hypersurface in every dimension ≥ 3 which is birational to a quartic del Pezzo surface fibration of type I_1 , and is stably rational over $\mathbb{Q}$.

Proposition 5.1. *The smooth cubic hypersurface*

$$X := \{(x_4 - 2x_3)x_1^2 + 3(x_4 + 2x_3)x_2^2 + 3x_3^2x_4 - x_4^3 + x_5^3 + \sum_{i=1}^r w_i^3 = 0\} \subset \mathbb{P}^{r+4}$$

is stably rational over $k = \mathbb{Q}$, for every $r \geq 0$.

Proof. When $r = 0$, one can directly check that X is smooth. Adding the Fermat cubic in variables w_i preserves the smoothness for every $r \geq 1$. Choose the line and the point

$$l = \{x_3 = x_4 = x_5 = w_1 = \dots = w_r = 0\}, \quad p = [0 : 0 : 0 : 1 : 0 : \dots : 0].$$

Projection from $\Pi = \langle l, p \rangle$ is given by

$$X \dashrightarrow \mathbb{P}^{r+1}, \quad (x_i, w_j) \mapsto (x_3, x_5, w_1, \dots, w_r).$$

Let

$$K = k(\mathbb{P}^{r+1}) = k(a, b_1, \dots, b_r).$$

We work on the affine chart $x_5 \neq 0$. Set $a = x_3/x_5$, and $b_i = w_i/x_5$. Renaming the coordinates

$$Y_1 = x_1, \quad Y_2 = x_2, \quad Y_3 = x_5, \quad Y_4 = x_4,$$

the generic fiber $X_\eta \subset \mathbb{P}_{Y_1, Y_2, Y_3, Y_4}^3$ is given by

$$Y_3(\beta Y_3^2 + 2a(3Y_2^2 - Y_1^2)) + Y_4(Y_1^2 + 3Y_2^2 + 3a^2Y_3^2 - Y_4^2) = 0, \quad (5.1)$$

where $\beta = 1 + \sum_{i=1}^r b_i^3$. One can check that X_η is smooth over K . Contracting the line

$$l_\eta = \{Y_3 = Y_4 = 0\} \subset \mathbb{P}^3,$$

we obtain a quartic del Pezzo surface $S \subset \mathbb{P}_{Y_1, \dots, Y_5}^4$ over K given by

$$\beta Y_3^2 + 2a(3Y_2^2 - Y_1^2) - Y_4Y_5 = Y_1^2 + 3Y_2^2 + 3a^2Y_3^2 - Y_4^2 + Y_3Y_5 = 0.$$

We show that S is stably rational over K . Consider the projection from the line l_η :

$$\pi_\eta : X_\eta \rightarrow \mathbb{P}^1, \quad (Y_1, Y_2, Y_3, Y_4) \mapsto (Y_3, Y_4).$$

Over the affine chart $t = Y_4/Y_3$, the generic fiber is the conic given by

$$f(Y_1, Y_2) := (t - 2a)Y_1^2 + 3(t + 2a)Y_2^2 - t^3 + 3a^2t + \beta = 0.$$

The discriminant of the quadratic form f is

$$\text{disc}(f) = 3(t - 2a)(t + 2a)(t^3 - 3a^2t - \beta).$$

Let ρ_1, ρ_2, ρ_3 be the roots of the irreducible polynomial

$$c(t) = t^3 - 3a^2t - \beta \in K[t].$$

A discriminant computation shows that

$$\Delta := \text{disc}(c(t)) = 27(2a^3 + \beta)(2a^3 - \beta)$$

is not a square in K . Thus $\text{Gal}(K_1/K) = \mathfrak{S}_3$, where $K_1 = K(\rho_1, \rho_2, \rho_3)$. The five singular fibers of π_η are individually defined over K_1 . Put

$$d_1 = \rho_2 - \rho_3, \quad d_2 = \rho_3 - \rho_1, \quad d_3 = \rho_1 - \rho_2.$$

The coefficients of $c(t)$ imply that

$$\rho_1 + \rho_2 + \rho_3 = 0, \quad \rho_1\rho_2 + \rho_2\rho_3 + \rho_1\rho_3 = -3a^2,$$

and thus

$$3(4a^2 - \rho_i^2) = d_i^2 \in K_1^{\times 2}, \quad i = 1, 2, 3.$$

It follows that singular fibers above ρ_1, ρ_2 , and ρ_3 split over K_1 . We label their components by

$$F_i^\pm : (\rho_i - 2a)Y_1 \pm d_iY_2 = 0. \quad (5.2)$$

On the other hand, singular fibers over $2a$ and $-2a$ split after adjoining e_1 and e_2 , respectively, where

$$e_1^2 = 3a(2a^3 - \beta), \quad e_2^2 = a(2a^3 + \beta).$$

We label their components by

$$F_4^\pm : 6aY_2 \pm e_1 = 0, \quad F_5^\pm : 2aY_1 \pm e_2 = 0. \quad (5.3)$$

Observe that $e_2 \in K_1(e_1)$. Indeed, one can check that

$$\left(\frac{e_1}{9(2a^3 - \beta)} \cdot d_1 d_2 d_3 \right)^2 = \left(\frac{e_1}{9(2a^3 - \beta)} \right)^2 \cdot \Delta = a(2a^3 + \beta).$$

All components of the singular fibers of π_η are individually defined over

$$K_1(e_1) = K(\rho_1, \rho_2, \rho_3, e_1),$$

which is thus the splitting field of X_η and S over K . There is a unique quadratic subextension $K(\sqrt{\Delta})/K$ of K_1/K . Assume that $e_1 \in K_1$. Then $K(\sqrt{\Delta}) = K(e_1)$, implying that either e_1^2 or e_1^2/Δ is a square in K . Computing their valuations at a , we see that this is impossible. Thus $e_1 \notin K_1$ and $K(e_1) \cap K_1 = K$. Since $K(e_1)K_1 = K_1(e_1)$, we know that

$$\text{Gal}(K_1(e_1)/K) = \text{Gal}(K(e_1)/K) \times \text{Gal}(K_1/K) = \mathfrak{C}_2 \times \mathfrak{S}_3.$$

Recall from Section 4 that the image of $\text{Gal}(K_1(e_1)/K)$ (or equivalently, the image of $\mathfrak{g}_K$) in $W(\mathbf{D}_5)$ is determined by its action on the 10 singular fibers (5.2) and (5.3).

- The (untwisted) 3-cycle from $\mathfrak{S}_3$ permutes ρ_i and thus d_i , and it fixes e_1 and e_2 . It maps to $(1, 2, 3) \in W(\mathbf{D}_5)$.
- Choose the (untwisted) 2-cycle τ from $\mathfrak{S}_3$ which fixes ρ_1 and e_1 , and switches ρ_2 and ρ_3 . Then $\tau(d_1) = -d_1$, $\tau(d_2) = -d_3$, $\tau(d_3) = -d_2$, and $\tau(e_2) = -e_2$. It maps to $c_1 c_2 c_3 c_5(2, 3)$.
- Let $\iota \in \text{Gal}(K_1(e_1)/K_1)$ be the order 2 element. We have that $\iota(d_i) = d_i$, for $i = 1, 2, 3$, and $\iota(e_i) = -e_i$, for $i = 1, 2$. It maps to $c_4 c_5 \in W(\mathbf{D}_5)$.

Therefore, the image of $\mathfrak{g}_K$ in $W(\mathbf{D}_5)$ is

$$\langle (1, 2, 3), c_1 c_2 c_3 c_5(2, 3), c_4 c_5 \rangle \subset W(\mathbf{D}_5),$$

which is conjugate to the group of type I_1 in Proposition 4.1. Applying Corollary 4.3, we conclude that S is stably rational over K . It follows that the smooth cubic X is stably rational over $k = \mathbb{Q}$. $\square$

The same arguments yield stably rational but nonrational smooth intersections of two quadrics in $\mathbb{P}^5$, over $\mathbb{Q}$ and over $\mathbb{C}(t)$. The following example was found using ChatGPT 5.6-Sol.

Example 5.2. Let $X = \{q_1 = q_2 = 0\} \subset \mathbb{P}^5$ be the smooth complete intersection of two quadrics, over $\mathbb{Q}$, where

$$\begin{aligned} q_1 &= 2x_1^2 - 6x_2^2 + 3x_3^2 - 3x_3x_4 + x_5^2 - x_4x_6, \\ q_2 &= x_1^2 + 3x_2^2 - x_4^2 - x_3x_6. \end{aligned}$$

Then

- (1) X contains no lines over $\mathbb{Q}$. In fact, X contains no lines over $\mathbb{R}$, since the quadratic form $q_1 - \frac{21}{10}q_2$ has signature $(1, 5)$. It follows that X is not rational over $\mathbb{Q}$ (and over $\mathbb{R}$), by [HT21] or [BW23].
- (2) The generic fiber of the rational map

$$f : X \dashrightarrow \mathbb{P}^1, \quad (x_1, \dots, x_6) \mapsto (x_3, x_5),$$

is a quartic del Pezzo surface S over $\mathbb{Q}(a)$ which is isomorphic to the surface (5.1) with $\beta = -3a^3 - a$. The same computation as above shows that S is of type I_1 , and thus X is stably rational, over $\mathbb{Q}$.

Type I_3 . Here, we construct a smooth cubic hypersurface over $\mathbb{Q}$, in every dimension ≥ 3 , that is birational to a quartic del Pezzo surface fibration of type I_3 , and is stably rational over $\mathbb{Q}$.

Proposition 5.3. *The smooth cubic hypersurface $X \subset \mathbb{P}^{r+4}$ given by*

$$x_4(x_1^2 + 2x_1x_2) + x_3(x_1^2 + x_1x_2 + x_2^2) + x_3^3 - x_3x_4^2 + x_4^3 + 2x_5^3 + \sum_{i=1}^r w_i^3 = 0$$

is stably rational over $k = \mathbb{Q}$, for any $r \geq 0$.

Proof. One can directly check that X is smooth when $r = 0$. Adding the Fermat cubic in variables w_i preserves the smoothness for every $r \geq 1$. Consider the projection

$$X \dashrightarrow \mathbb{P}^{r+1}, \quad (x_i, w_j) \rightarrow (x_4, x_5, w_1, \dots, w_r).$$

Let $K = k(\mathbb{P}^{r+1}) = k(a, b_1, \dots, b_r)$. Set $a = x_4/x_5$, and $b_i = w_i/x_5$. Renaming the coordinates

$$Y_1 = x_1, \quad Y_2 = x_2, \quad Y_3 = x_5, \quad Y_4 = x_3,$$

the generic fiber $X_\eta \subset \mathbb{P}_{Y_1, Y_2, Y_3, Y_4}^3$ is given by

$$Y_3(aY_1^2 + 2aY_1Y_2 + (a^3 + \beta)Y_3^2) + Y_4(Y_1^2 + Y_1Y_2 + Y_2^2 - a^2Y_3^2 + Y_4^2) = 0,$$

where $\beta = 2 + \sum_{i=1}^r b_i^3$. One can check that X_η is smooth over K . Contracting the line $l_\eta = \{Y_3 = Y_4 = 0\} \subset \mathbb{P}^3$ yields a quartic del Pezzo surface S birational to X_η over K . Consider the conic bundle obtained via projection from l_η

$$\pi_\eta : X_\eta \rightarrow \mathbb{P}^1, \quad (Y_1, Y_2, Y_3, Y_4) \rightarrow (Y_3, Y_4).$$

Over the affine chart $t = Y_4/Y_3$, the generic fiber is the conic given by

$$f(Y_1, Y_2) := (t + a)Y_1^2 + (t + 2a)Y_1Y_2 + tY_2^2 + t^3 - a^2t + a^3 + \beta = 0.$$

The discriminant of the quadratic form f is

$$\text{disc}(f) = -\frac{1}{4}(t^3 - a^2t + a^3 + \beta)(3t^2 - 4a^2).$$

Let $c_1(t) = t^3 - a^2t + a^3 + \beta$, and ρ_1, ρ_2, ρ_3 its three roots. One sees that

$$\Delta := \text{disc}(c_1) = -23a^6 - 54a^3\beta - 27\beta^2$$

is not a square in K . Thus $\text{Gal}(K_1/K) = \mathfrak{S}_3$, where $K_1 = K(\rho_1, \rho_2, \rho_3)$. Put

$$d_1 = \rho_2 - \rho_3, \quad d_2 = \rho_3 - \rho_1, \quad d_3 = \rho_1 - \rho_2.$$

The coefficients of $c_1(t)$ imply that

$$d_i^2 = 4a^2 - 3\rho_i^2.$$

The singular fibers over ρ_i split over K_1 , and their components are

$$F_i^\pm : 2(\rho_i + a)Y_1 + (\rho_i + 2a \pm d_i)Y_2 = 0, \quad i = 1, 2, 3. \quad (5.4)$$

Let $c_2(t) = 3t^2 - 4a^2$. The splitting field of c_2 is $K(\sqrt{3})$, and c_2 has two roots $\rho_4 = \frac{2a}{\sqrt{3}}$ and $\rho_5 = -\frac{2a}{\sqrt{3}}$. We check that $\sqrt{3} \notin K_1$. Indeed, $K(\sqrt{3})$ is different from the unique quadratic subextension $K(\sqrt{\Delta})/K$ of K_1/K , since $\Delta/3$ is not a square in K . Set $K_2 = K_1(\sqrt{3})$. It follows that

$$\text{Gal}(K_2/K) = \text{Gal}(K(\sqrt{3})/K) \times \text{Gal}(K_1/K) = \mathfrak{C}_2 \times \mathfrak{S}_3.$$

Over K_2 , each singular fiber of π_η is defined. Choose $D_1 \in \bar{K}$ such that

$$D_1^2 = (-32\sqrt{3} - 52)a^4 - (24\sqrt{3} + 36)a\beta,$$

and put

$$D_2 = \frac{4a}{D_1}d_1d_2d_3. \quad (5.5)$$

One can check that

$$D_2^2 = (32\sqrt{3} - 52)a^4 + (24\sqrt{3} - 36)a\beta.$$

The singular fibers over ρ_4 and ρ_5 split over $K_2(D_1)$, with components:

$$\begin{aligned} F_4^\pm : 2a(3 + 2\sqrt{3})Y_1 + a(6 + 2\sqrt{3})Y_2 \pm D_1 &= 0, \\ F_5^\pm : 2a(3 - 2\sqrt{3})Y_1 + a(6 - 2\sqrt{3})Y_2 \pm D_2 &= 0. \end{aligned} \quad (5.6)$$

Thus, the splitting field of X_η and S is

$$K_2(D_1) = K(\sqrt{3}, \rho_1, \rho_2, \rho_3)(D_1) = K_1(D_1).$$

We show that $D_1 \notin K_2$. One can check that the minimal polynomial of D_1 over K is

$$(t^2 - D_1^2)(t^2 - D_2^2) \in K[t]. \quad (5.7)$$

Then $K(D_1)/K$ has degree 4 and is not Galois since $D_2 \notin K(D_1)$. However, any index 4 subgroup in $\text{Gal}(K_2/K) = \mathfrak{S}_2 \times \mathfrak{S}_3$ is normal, which implies that $D_1 \notin K_2$. It follows that

$$[K_1(D_1) : K] = [K_2(D_1) : K] = [K_2(D_1) : K_2][K_2 : K] = 24.$$

Let $G = \text{Gal}(K_1(D_1)/K)$. From (5.5), one sees that $K_2(D_1)$ is the splitting field of (5.7) over K_1 . It follows that $K_1(D_1)/K_1$ is Galois, and $\text{Gal}(K_1(D_1)/K_1) = \mathfrak{S}_2^2$, since it has 3 different quadratic subextensions $K_1(\sqrt{3})$, $K_1(D_1 + D_2)$ and $K_1(D_1 - D_2)$. We determine the image of $\text{Gal}(K_1(D_1)/K_1)$ in $W(\mathbf{D}_5)$ by identifying its action on (5.6).

- One generator of $\text{Gal}(K_1(D_1)/K_1)$ swaps D_1 and D_2 , and maps to $(4, 5) \in W(\mathbf{D}_5)$.
- The other generator of $\text{Gal}(K_1(D_1)/K_1)$ changes the signs of D_1 and D_2 simultaneously, and maps to $c_4 c_5 \in W(\mathbf{D}_5)$.

The subextension

$$K_1(D_1) \supset K_1 \supset K$$

yields an extension of Galois groups

$$1 \rightarrow \mathfrak{S}_2^2 \rightarrow G \rightarrow \mathfrak{S}_3 \rightarrow 1. \quad (5.8)$$

The extension (5.8) splits since we know that $K_1 \cap K(D_1) = K$. Any element in $\text{Gal}(K_1/K) = \mathfrak{S}_3$ can be lifted to G as an element fixing D_1 . We describe their actions on singular fibers (5.4) and (5.6).

- A lift of $(1, 2, 3) \in \mathfrak{S}_3$ to G permutes $\rho_1 \mapsto \rho_2 \mapsto \rho_3$, and fixes D_1 and D_2 . Its image in $W(\mathbf{D}_5)$ is $(1, 2, 3)$.
- A lift τ of $(2, 3) \in \mathfrak{S}_3$ fixes ρ_1 and D_1 , and swaps ρ_2 and ρ_3 . One can check that $\tau(d_1) = -d_1$, $\tau(d_2) = -d_3$, $\tau(d_3) = -d_2$, and thus $\tau(D_2) = -D_2$. Its image in $W(\mathbf{D}_5)$ is $c_1 c_2 c_3 c_5(2, 3)$.

Thus the image of $\mathfrak{g}_K$ in $W(\mathbf{D}_5)$ is

$$\langle (4, 5), c_4 c_5, (1, 2, 3), c_1 c_2 c_3 c_5(2, 3) \rangle,$$

which is conjugate to the group of type I_3 listed in Proposition 4.1. Applying Corollary 4.3, we conclude that X is stably rational over $\mathbb{Q}$. $\square$

Remark 5.4. The cubic hypersurfaces in Propositions 5.1 and 5.3 are not isomorphic over $\mathbb{C}$. One can see this by checking that the cubic surfaces $X \cap \{x_5 = w_1 = \dots = w_r = 0\}$ in the two cases have different numbers of Eckardt points.

It would be interesting to study quartic del Pezzo fibration structures of types I_0, I_1, I_2, I_3 on other nonrational varieties, such as singular quartic hypersurfaces or quartic double solids, cf. [Che06], [Voi15].

6. STABLE LINEARIZABILITY

Recall that a generically free regular G -action on a variety X is called stably linearizable if $X \times \mathbb{P}^r$ is equivariantly birational to the projectivization of a linear G -representation, with the trivial G -action on $\mathbb{P}^r$, for some $r \geq 0$. Here, we give a detailed proof of stable linearizability of the unique G -minimal nonlinearizable del Pezzo surface satisfying Condition **(H1)**, see [Pro15, Theorem 1.2]. We work over a field k of characteristic zero containing ζ_3 , a primitive third root of 1.

Proposition 6.1. *Let S be the del Pezzo surface of degree 4 given by*

$$S = \{x_1^2 + \zeta_3 x_2^2 + \zeta_3^2 x_3^2 + x_4^2 = x_1^2 + \zeta_3^2 x_2^2 + \zeta_3 x_3^2 + x_5^2 = 0\},$$

with an action of $G = \mathfrak{S}_3 \rtimes \mathfrak{S}_4$ generated by

$$\begin{aligned} \sigma : (x) &\mapsto (x_2, x_3, x_1, \zeta_3 x_4, \zeta_3^2 x_5), \\ \tau : (x) &\mapsto (x_1, x_3, x_2, -x_5, x_4). \end{aligned}$$

Then the G -action on S is stably linearizable.

A quick proof of this proposition, based on [DR15], would go as follows: a G -variety is stably linearizable if and only if all twists (or equivalently, one *versal* twist) are stably rational, see [DR15, Theorem 1.1], combined with [KT26b, Theorem 2.5]. The key is that $S^G \neq \emptyset$, so that for every twist over a field K/k , the set of K -points of a form of S is nonempty. Then Corollary 1.2 gives stable rationality over K . Nevertheless, we include a direct proof within equivariant geometry, based on the equivariant version of the torsor formalism developed in [HT23] and [KT25].

Proof of Proposition 6.1. Note that k is the splitting field of S . Thus, over k , there is a unique universal torsor over S . By [HT23, Corollary 9], it suffices to show that

- (1) $\text{Pic}(S)$ is a G -stably permutation module;
- (2) the G -action lifts to the universal torsor $\mathcal{T}$ of S ;
- (3) the G -action on $\mathcal{T}$ is linearizable.

We have proved (1) in Lemma 4.2. By [HT23], (2) follows from the fact that $S^G \neq \emptyset$. To show (3), we repeat the Cox ring computation from Section 3 with the same notation, noting that S is the blowup of $\mathbb{P}^2$ at the following 5 points:

$$p_1 = [1 : 0 : 0], \quad p_2 = [0 : 1 : 0], \quad p_3 = [0 : 0 : 1], \quad p_4 = [1 : 1 : 1],$$

$$p_5 = [1 : \zeta_3 : \zeta_3^2].$$

In the anticanonical model, the exceptional curves can be identified as follows

$$\begin{aligned}
E_1 &= \{x_3 - x_1 + x_2 = x_4 - \zeta_3^2 x_1 - x_2 = x_5 - \zeta_3 x_1 - x_2 = 0\}, \\
E_2 &= \{x_3 + x_1 - x_2 = x_4 + \zeta_3^2 x_1 + x_2 = x_5 + \zeta_3 x_1 + x_2 = 0\}, \\
E_3 &= \{x_3 - x_1 - x_2 = x_4 + \zeta_3^2 x_1 - x_2 = x_5 + \zeta_3 x_1 - x_2 = 0\}, \\
E_4 &= \{x_3 + x_1 + x_2 = x_4 - \zeta_3^2 x_1 + x_2 = x_5 + \zeta_3 x_1 - x_2 = 0\}, \\
E_5 &= \{x_3 + x_1 + x_2 = x_4 + \zeta_3^2 x_1 - x_2 = x_5 - \zeta_3 x_1 + x_2 = 0\}, \\
L_{12} &= \{x_3 - x_1 - x_2 = x_4 - \zeta_3^2 x_1 + x_2 = x_5 - \zeta_3 x_1 + x_2 = 0\}, \\
L_{13} &= \{x_3 + x_1 - x_2 = x_4 - \zeta_3^2 x_1 - x_2 = x_5 - \zeta_3 x_1 - x_2 = 0\}, \\
L_{14} &= \{x_3 - x_1 + x_2 = x_4 + \zeta_3^2 x_1 + x_2 = x_5 - \zeta_3 x_1 - x_2 = 0\}, \\
L_{15} &= \{x_3 - x_1 + x_2 = x_4 - \zeta_3^2 x_1 - x_2 = x_5 + \zeta_3 x_1 + x_2 = 0\}, \\
L_{23} &= \{x_3 - x_1 + x_2 = x_4 + \zeta_3^2 x_1 + x_2 = x_5 + \zeta_3 x_1 + x_2 = 0\}, \\
L_{24} &= \{x_3 + x_1 - x_2 = x_4 - \zeta_3^2 x_1 - x_2 = x_5 + \zeta_3 x_1 + x_2 = 0\}, \\
L_{25} &= \{x_3 + x_1 - x_2 = x_4 + \zeta_3^2 x_1 + x_2 = x_5 - \zeta_3 x_1 - x_2 = 0\}, \\
L_{34} &= \{x_3 - x_1 - x_2 = x_4 - \zeta_3^2 x_1 + x_2 = x_5 + \zeta_3 x_1 - x_2 = 0\}, \\
L_{35} &= \{x_3 - x_1 - x_2 = x_4 + \zeta_3^2 x_1 - x_2 = x_5 - \zeta_3 x_1 + x_2 = 0\}, \\
L_{45} &= \{x_3 + x_1 + x_2 = x_4 - \zeta_3^2 x_1 + x_2 = x_5 - \zeta_3 x_1 + x_2 = 0\}, \\
Q &= \{x_3 + x_1 + x_2 = x_4 + \zeta_3^2 x_1 - x_2 = x_5 + \zeta_3 x_1 - x_2 = 0\}.
\end{aligned}$$

We present an explicit lift of the G -action to $\text{Cox}(S)$, and verify that it is compatible with the G -action on S . Consider the G -action given by

$$\begin{aligned}
\sigma(q) &= q, & \sigma(e_1) &= e_2, & \sigma(e_2) &= e_3, & \sigma(e_3) &= e_1, \\
\sigma(e_4) &= e_4, & \sigma(e_5) &= e_5, & \sigma(\ell_{12}) &= \zeta_3 \ell_{23}, & \sigma(\ell_{13}) &= \zeta_3 \ell_{12}, \\
\sigma(\ell_{14}) &= -\zeta_3 \ell_{24}, & \sigma(\ell_{15}) &= -\ell_{25}, & \sigma(\ell_{23}) &= \zeta_3 \ell_{13}, & \sigma(\ell_{24}) &= -\zeta_3 \ell_{34}, \\
\sigma(\ell_{25}) &= -\ell_{35}, & \sigma(\ell_{34}) &= \zeta_3 \ell_{14}, & \sigma(\ell_{35}) &= \ell_{15}, & \sigma(\ell_{45}) &= \ell_{45},
\end{aligned}$$

and

$$\begin{aligned}
\tau(q) &= 3(\zeta_3^2 - 1)e_4, & \tau(e_1) &= \frac{\zeta_3^2 - 1}{3}\ell_{14}, & \tau(e_2) &= \frac{\zeta_3 - \zeta_3^2}{3}\ell_{34}, \\
\tau(e_3) &= \frac{\zeta_3 - 1}{3}\ell_{24}, & \tau(e_4) &= \frac{\zeta_3^2 - 1}{9}\ell_{45}, & \tau(e_5) &= \frac{\zeta_3 - 1}{9}q, \\
\tau(\ell_{12}) &= \frac{\zeta_3^2 - 1}{3}\ell_{25}, & \tau(\ell_{13}) &= \frac{\zeta_3 - 1}{3}\ell_{35}, & \tau(\ell_{14}) &= (\zeta_3^2 - \zeta_3)\ell_{23}, \\
\tau(\ell_{15}) &= (1 - \zeta_3)e_1, & \tau(\ell_{23}) &= \frac{\zeta_3^2 - \zeta_3}{3}\ell_{15}, & \tau(\ell_{24}) &= (\zeta_3^2 - 1)\ell_{12}, \\
\tau(\ell_{25}) &= (\zeta_3 - 1)e_3, & \tau(\ell_{34}) &= (\zeta_3 - 1)\ell_{13}, & \tau(\ell_{35}) &= (1 - \zeta_3)e_2, \\
\tau(\ell_{45}) &= 3(\zeta_3 - 1)e_5.
\end{aligned}$$

One can check that the ideal $\langle R_1, \dots, R_{20} \rangle$ given in Section 3 is G -invariant, and the permutation action of G on the generators agree with that on the exceptional curves listed above. In particular, the G -action on $\text{Cox}(S)$ respects the grading by $\text{Pic}(S)$. Under the following basis of $H^0(S, -K_S)$:

$$\begin{aligned}
s_1 &= qe_2e_3\ell_{23} - e_1\ell_{12}\ell_{13}\ell_{45}, \\
s_2 &= \zeta_3qe_1e_3\ell_{13} - \zeta_3^2e_2\ell_{12}\ell_{23}\ell_{45}, \\
s_3 &= qe_2e_3\ell_{23} + \zeta_3qe_1e_3\ell_{13} + 2\zeta_3^2qe_1e_2\ell_{12} + \zeta_3^2e_2\ell_{12}\ell_{23}\ell_{45} + e_1\ell_{12}\ell_{13}\ell_{45}, \\
s_4 &= \zeta_3^2qe_2e_3\ell_{23} - \zeta_3qe_1e_3\ell_{13} - \zeta_3^2e_2\ell_{12}\ell_{23}\ell_{45} + \zeta_3^2e_1\ell_{12}\ell_{13}\ell_{45}, \\
s_5 &= \zeta_3qe_2e_3\ell_{23} - \zeta_3qe_1e_3\ell_{13} - \zeta_3^2e_2\ell_{12}\ell_{23}\ell_{45} + \zeta_3e_1\ell_{12}\ell_{13}\ell_{45},
\end{aligned}$$

the resulting anticanonical map

$$\pi : \text{Spec}(\text{Cox}(S)) \xrightarrow{|-K_S|} S$$

is equivariant under the prescribed G -action given in Proposition 6.1.

To show the linearizability of the G -action on $\mathcal{T}$, it suffices to establish that for the G -action on $\text{Spec}(\text{Cox}(S))$. The fixed locus $\mathbb{P}(U)^G$ consists of four points. Consider the projection from the affine tangent space of one of these points, given by

$$\varphi : \text{Spec}(\text{Cox}(S)) \dashrightarrow \mathbb{A}^8, \quad (q, e_1, \dots, e_5, l_{ij}) \mapsto (f_1, \dots, f_8)$$

where,

$$\begin{aligned}
f_1 &= e_2 - \zeta_3 e_3 + (\zeta_3 - 1)e_4 + l_{12} - l_{13} + l_{14}, \\
f_2 &= \zeta_3 e_2 - e_3 - (\zeta_3 - 1)e_5 + \zeta_3 l_{12} - \zeta_3^2 l_{13} + l_{15}, \\
f_3 &= e_1 - \zeta_3^2 e_3 - (\zeta_3 + 2)e_4 + l_{12} - l_{23} + l_{24}, \\
f_4 &= e_1 - \zeta_3 e_3 + (\zeta_3 - 1)e_5 + l_{12} - \zeta_3^2 l_{23} + l_{25}, \\
f_5 &= \zeta_3 e_1 - \zeta_3^2 e_2 - (2\zeta_3 + 1)e_4 + l_{13} - l_{23} + l_{34}, \\
f_6 &= \zeta_3 e_1 - e_2 - (\zeta_3 - 1)e_5 + l_{13} - \zeta_3 l_{23} + l_{35}, \\
f_7 &= 3e_1 - 3\zeta_3 e_2 + 3(\zeta_3 - 1)e_5 - (\zeta_3 + 2)l_{14} + (2\zeta_3 + 1)l_{24} + l_{45}, \\
f_8 &= q - 3(\zeta_3 + 2)e_4 + 3l_{12} - 3\zeta_3 l_{13} - (2\zeta_3 + 1)l_{25} + (\zeta_3 + 2)l_{35}.
\end{aligned}$$

An explicit inverse of φ , given in [TZ26], proves that φ is birational. $\square$

7. ARITHMETIC APPLICATIONS

J.-L. Colliot-Thélène encouraged us to include the following consequences of Theorem 3.4, i.e., the rationality of a universal torsor $\mathcal{T}$ over a quartic del Pezzo surface S , provided $\mathcal{T}(k) \neq \emptyset$:

- (1) The evaluation map from $S(k)/R$ (the set of rational points modulo R -equivalence) to $H^1(k, T)$ is injective.
- (2) Let $A_0(X) \subset \text{CH}_0(X)$ be the Chow group of zero-cycles of degree zero modulo rational equivalence, and $p_0 \in S(k)$. Then the map

$$S(k)/R \rightarrow A_0(S), \quad p \mapsto p - p_0$$

is bijective. Its surjectivity was proved in [CTC79].

- (3) Let k be a finitely generated extension of $\mathbb{Q}$. Then $S(k)/R$ is finite. This follows from (2) and the finiteness of $A_0(X)$, proved in [CTS81] and [Blo81].

These were known when S admits a conic bundle structure, by [CTS87b].

REFERENCES

- [BCTSSD85] A. Beauville, J.-L. Colliot-Thélène, J.-J. Sansuc, and P. Swinnerton-Dyer. Variétés stablement rationnelles non rationnelles. *Ann. of Math.* (2), 121(2):283–318, 1985.
- [Blo81] S. Bloch. On the Chow groups of certain rational surfaces. *Ann. Sci. École Norm. Sup. (4)*, 14(1):41–59, 1981.
- [BP04] V. V. Batyrev and O. N. Popov. The Cox ring of a del Pezzo surface. In *Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002)*, volume 226 of *Progr. Math.*, pages 85–103. Birkhäuser, 2004.
- [BW23] O. Benoist and O. Wittenberg. Intermediate Jacobians and rationality over arbitrary fields. *Ann. Sci. Éc. Norm. Supér. (4)*, 56(4):1029–1084, 2023.

- [CG72] C. H. Clemens and P. A. Griffiths. The intermediate Jacobian of the cubic threefold. *Ann. of Math. (2)*, 95:281–356, 1972.
- [Che06] I. Cheltsov. Nonrational nodal quartic threefolds. *Pacific J. Math.*, 226(1):65–81, 2006.
- [CMR04] C. Ciliberto, M. Mella, and F. Russo. Varieties with one apparent double point. *J. Algebraic Geom.*, 13(3):475–512, 2004.
- [CR06] C. Ciliberto and F. Russo. Varieties with minimal secant degree and linear systems of maximal dimension on surfaces. *Adv. Math.*, 200(1):1–50, 2006.
- [CTC79] J.-L. Colliot-Thélène and D. Coray. L'équivalence rationnelle sur les points fermés des surfaces rationnelles fibrées en coniques. *Compositio Math.*, 39(3):301–332, 1979.
- [CTS77] J.-L. Colliot-Thélène and J.-J. Sansuc. La R -équivalence sur les tores. *Ann. Sci. École Norm. Sup. (4)*, 10(2):175–229, 1977.
- [CTS81] J.-L. Colliot-Thélène and J.-J. Sansuc. On the Chow groups of certain rational surfaces: a sequel to a paper of S. Bloch. *Duke Math. J.*, 48(2):421–447, 1981.
- [CTS87a] J.-L. Colliot-Thélène and J.-J. Sansuc. La descente sur les variétés rationnelles. II. *Duke Math. J.*, 54(2):375–492, 1987.
- [CTS87b] J.-L. Colliot-Thélène and A. N. Skorobogatov. R -equivalence on conic bundles of degree 4. *Duke Math. J.*, 54(2):671–677, 1987.
- [Der07] U. Derenthal. Universal torsors of del Pezzo surfaces and homogeneous spaces. *Adv. Math.*, 213(2):849–864, 2007.
- [DP19] U. Derenthal and M. Pieropan. Cox rings over nonclosed fields. *J. Lond. Math. Soc. (2)*, 99(2):447–476, 2019.
- [DR15] A. Duncan and Z. Reichstein. Versality of algebraic group actions and rational points on twisted varieties. *J. Algebr. Geom.*, 24(3):499–530, 2015.
- [EFS25] P. Engel, O.d.G. Fortman, and S. Schreieder. Matroids and the integral Hodge conjecture for abelian varieties, 2025. [arXiv:2507.15704](#).
- [HKT22] B. Hassett, J. Kollár, and Yu. Tschinkel. Rationality of even-dimensional intersections of two real quadrics. *Comment. Math. Helv.*, 97(1):183–207, 2022.
- [HKT23] B. Hassett, A. Kresch, and Yu. Tschinkel. Stable rationality in smooth families of threefolds. *Duke Math. J.*, 172(6):1145–1172, 2023.
- [HT17a] B. Hassett and Yu. Tschinkel. Stable rationality of quartic del Pezzo surfaces, 2017. unpublished note.
- [HT17b] B. Hassett and Yu. Tschinkel. Stably rational cubic threefolds, 2017. unpublished note.
- [HT21] B. Hassett and Yu. Tschinkel. Rationality of complete intersections of two quadrics over nonclosed fields. *Enseign. Math.*, 67(1-2):1–44, 2021. With an appendix by Jean-Louis Colliot-Thélène.
- [HT23] B. Hassett and Yu. Tschinkel. Torsors and stable equivariant birational geometry. *Nagoya Math. J.*, 250:275–297, 2023.
- [KiST89] B. Kunyavskiĭ, A. Skorobogatov, and M. Tsfasman. del Pezzo surfaces of degree four. *Mém. Soc. Math. France (N.S.)*, (37):113, 1989.
- [KT25] A. Kresch and Yu. Tschinkel. Equivariant unirationality of toric varieties, 2025. [arXiv:2506.07152](#).

- [KT26a] A. Kresch and Yu. Tschinkel. Invariants in equivariant birational geometry, 2026. [arXiv:2602.23998](#).
- [KT26b] A. Kresch and Yu. Tschinkel. Linearizability notions in equivariant birational geometry, 2026. [arXiv:2606.10965](#).
- [Man66] Yu. I. Manin. Rational surfaces over perfect fields. *Inst. Hautes Études Sci. Publ. Math.*, (30):55–113, 1966.
- [Man86] Yu. I. Manin. *Cubic forms*, volume 4 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, second edition, 1986.
- [Pro15] Yu. Prokhorov. On stable conjugacy of finite subgroups of the plane Cremona group, II. *Michigan Math. J.*, 64(2):293–318, 2015.
- [SB04] N. I. Shepherd-Barron. Stably rational irrational varieties. In *The Fano Conference*, pages 693–700. Univ. Torino, Turin, 2004.
- [SS91] P. Salberger and A. N. Skorobogatov. Weak approximation for surfaces defined by two quadratic forms. *Duke Math. J.*, 63(2):517–536, 1991.
- [SS07] V. Serganova and A. Skorobogatov. Del Pezzo surfaces and representation theory. *Algebra Number Theory*, 1(4):393–419, 2007.
- [TY20] Yu. Tschinkel and K. Yang. Potentially stably rational del Pezzo surfaces over nonclosed fields. In *Combinatorial and additive number theory. III*, volume 297 of *Springer Proc. Math. Stat.*, pages 227–233. Springer, Cham, 2020.
- [TZ26] Yu. Tschinkel and Zh. Zhang. Supporting materials, 2026. available at: <https://zhijiazhangz.github.io/dP4/>.
- [Voi15] C. Voisin. Unirational threefolds with no universal codimension 2 cycle. *Invent. Math.*, 201(1):207–237, 2015.
- [Voi17] C. Voisin. On the universal CH_0 group of cubic hypersurfaces. *J. Eur. Math. Soc. (JEMS)*, 19(6):1619–1653, 2017.

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA

SIMONS FOUNDATION, 160 FIFTH AVENUE, NEW YORK, NY 10010, USA

Email address: tschinkel@cims.nyu.edu

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027, USA

Email address: zz3531@columbia.edu